

IMPROVEMENT TAYKOV'S LOWER BOUND IN AN INEQUATION BETWEEN C AND L NORMS FOR TRIGONOMETRIC POLYNOMIALS

D.V. GORBACHEV

ABSTRACT. We give new lower asymptotical estimate of constant

$$C_n = \sup \left\{ \frac{\|t_n\|_{C(\mathbb{T})}}{\|t_n\|_{L(\mathbb{T})}} : t_n \text{ are real trigonometric polynomials, } \deg t_n < n \right\}$$

as $n \rightarrow \infty$. This estimate improves known bound of L.V.Taykov (1965).

Suppose $e(t) = \exp(2\pi it)$, $c(t) = \cos(2\pi t)$, $s(t) = \sin(2\pi t)$. We consider an extremum problem about behavior of the best constant as $n \rightarrow \infty$ in the inequation between C and L norms for real trigonometric polynomials $t_n(x) = \sum_{\nu \in \mathbb{Z}, |\nu| < n} \hat{t}_n(\nu) e(\nu x)$:

$$C_n = \sup_{t_n} \frac{\|t_n\|_C}{\|t_n\|_L},$$

where

$$\|t_n\|_C = \max_{|x| \leq 1/2} |t_n(x)|, \quad \|t_n\|_L = \int_{-1/2}^{1/2} |t_n(x)| dx.$$

Exact value of the constant C_n is unknown for all $n \geq 3$ [1] ($C_1 = 1$, $C_2 = 2,12532 \dots$). S.B. Stechkin (see [1, 2]) proved that $C_n = c \cdot n + o(n)$ as $n \rightarrow \infty$ ($c > 0$). L.V. Taykov [2] improved this result. He found following bounds for C_n :

$$(1,07995 \dots) n + O(1) = \frac{2}{\int_0^\pi \sin t dt/t} n + O(1) \leq C_n \leq \frac{4G}{\pi} n + O(1) = (1,16624 \dots) n + O(1), \quad (1)$$

where $G = \sum_{k=0}^{\infty} (-1)^k (2k+1)^{-2} = 0,91596 \dots$ is Katalan's constant. The lower bound of this result is obtained for polynomials of Rogozinsky [3]

$$R_n(x) = \sum_{|\nu| < n} c\left(\frac{\nu}{4n}\right) e(\nu x) = s\left(\frac{1}{4n}\right) c(nx) (c(x) - c\left(\frac{1}{4n}\right))^{-1}. \quad (2)$$

In this work we show new polynomials for improvement the lower bound in (1). By $q_k = k/2 - 1/4$ ($k \in \mathbb{Z}$) denote zeros of function $c(t)$; $1/2 < z_1 < 1 < z_2 < 3/2$,

$$\omega(z) = \frac{(z_1^2 - z^2)(z_2^2 - z^2)}{(q_1^2 - z^2)(q_2^2 - z^2)(q_3^2 - z^2)}, \quad \omega_k(z) = (q_k^2 - z^2)\omega(z) \quad (k = 1, 2, 3).$$

Consider an even function

$$\varphi(x) = \sum_{k=1}^3 \omega_k(q_k) \sigma_k(x)$$

such that $\sigma_k(x) = (\pi/q_k) s(q_k(1 - |x|))$ for $|x| < 1$ and $\sigma_k(x) = 0$ for $|x| \geq 1$.

Fourier transform $\hat{\sigma}_k(z) = \int_{\mathbb{R}} \sigma_k(x) e(-zx) dx$ equals to $c(z)(q_k^2 - z^2)^{-1}$. Therefore Fourier transform of the function φ equals to

$$\hat{\varphi}(z) = \sum_{k=1}^3 \omega_k(q_k) \frac{c(z)}{q_k^2 - z^2} = \omega(z) c(z).$$

Suppose

$$F_n(x) = \sum_{|\nu| < n} \varphi\left(\frac{\nu}{n}\right) e(\nu x).$$

Using Poisson summation formula $\sum_{\nu \in \mathbb{Z}} \hat{f}(\nu - x) = \sum_{\nu \in \mathbb{Z}} f(\nu) e(\nu x)$ for the function $f(x) = \varphi(x/n)$ (since $\hat{\varphi}(z) = O(z^{-2})$ ($|z| \rightarrow \infty$), $\varphi(x) = 0$ ($|x| \geq 1$)) and identity

$$\sum_{\nu \in \mathbb{Z}} \frac{1}{a^2 - (z + \nu)^2} = \frac{(\pi/a)s(a)}{c(z) - c(a)},$$

we get a following expression for polynomial F_n

$$F_n(x) = n \sum_{\nu \in \mathbb{Z}} \hat{\varphi}(nx + n\nu) = c(nx) \sum_{k=1}^3 \frac{(\pi/q_k)\omega_k(q_k)s(\frac{q_k}{n})}{c(x) - c(\frac{q_k}{n})}. \quad (3)$$

THEOREM. $\|F_n\|_C / \|F_n\|_L \geq \rho(\alpha, \beta) n + O(n^{-1})$, where

$$\rho(z_1, z_2) = \hat{\varphi}(0) \left[4 \left(\int_0^{q_1} + \int_{q_2}^{q_3} - \int_{z_1}^{z_2} \right) \hat{\varphi}(z) dz \right]^{-1}.$$

The function $\rho(z_1, z_2)$ can be expressed by an integral sine. Using necessary conditions of extreme, we maximize this function on $1/2 < z_1 < 1 < z_2 < 3/2$ and obtain new asymptotic lower bound for the constant C_n .

COROLLARY. $C_n \geq (1.08176 \dots) n + O(n^{-1})$, $z_1^* = 0.72096 \dots$, $z_2^* = 1.23305 \dots$

REMARK 1. For $z_1 = q_2$, $z_2 = q_3$ polynomials F_n coincide with polynomials of Rogozinsky (2) (accurate to a coefficient 4π) $F_n(x) = 4\pi R_n(x)$ and $\rho(q_2, q_3) = 2(\int_0^\pi \sin t dt/t)^{-1}$; this fact give Taykov's lower bound.

PROOF. Since $\hat{\varphi}(z) = O(z^{-2})$ as $|z| \rightarrow \infty$, it follows that $n \sum_{\nu \in \mathbb{Z} \setminus \{0\}} \hat{\varphi}(nx + n\nu) = O(n^{-1})$ uniformly for $|x| \leq 1/2$. Therefore from (3) we have

$$\|F_n\|_C = \max_{|x| \leq 1/2} \left| n \sum_{\nu \in \mathbb{Z}} \hat{\varphi}(nx + n\nu) \right| = \max_{|z| \leq n/2} |n\hat{\varphi}(z)| + O(n^{-1}) \geq n\hat{\varphi}(0) + O(n^{-1}).$$

Now we estimate $\|F_n\|_L$. We have

$$\begin{aligned} \|F_n\|_L &= \int_{-1/2}^{1/2} \left| n \sum_{\nu \in \mathbb{Z}} \hat{\varphi}(nx + n\nu) \right| dx = \int_{-1/2}^{1/2} |n\hat{\varphi}(nx)| dx + O(n^{-1}) = \\ &= \int_{-n/2}^{n/2} |\hat{\varphi}(z)| dz + O(n^{-1}) \leq \int_{\mathbb{R}} |\hat{\varphi}(z)| dz + O(n^{-1}). \end{aligned}$$

Therefore $\|F_n\|_C / \|F_n\|_L \geq \frac{\hat{\varphi}(0)}{\int_{\mathbb{R}} |\hat{\varphi}(z)| dz} n + O(n^{-1})$.

We calculate $\int_{\mathbb{R}} |\hat{\varphi}(z)| dz$. Suppose

$$E_+ = [0, z_1] \cup [z_2, q_4] \cup (\cup_{k=2}^\infty [q_{2k+1}, q_{2k+2}]), \quad E_- = (z_1, z_2) \cup (\cup_{k=2}^\infty (q_{2k}, q_{2k+1})).$$

Then $\mathbb{R}_+ = E_+ \cup E_-$ and $\hat{\varphi}(z) \geq 0$ for $z \in E_+$; $\hat{\varphi}(z) < 0$ for $z \in E_-$. $\hat{\varphi}$ is even function. Hence we obtain

$$\begin{aligned} \int_{\mathbb{R}} |\hat{\varphi}(z)| dz &= 2 \int_{\mathbb{R}_+} |\hat{\varphi}(z)| dz = 2 \left(\int_0^{z_1} + \int_{z_2}^{q_4} + \sum_{k=2}^\infty \int_{q_{2k+1}}^{q_{2k+2}} - \int_{z_1}^{z_2} - \sum_{k=2}^\infty \int_{q_{2k}}^{q_{2k+1}} \right) \hat{\varphi}(z) dz = \\ &= 2 \left(\int_{\mathbb{R}_+} -2 \left(\int_{z_1}^{z_2} + \sum_{k=2}^\infty \int_{k-1/4}^{k+1/4} \right) \right) \hat{\varphi}(z) dz = \varphi(0) - 4 \int_{z_1}^{z_2} \hat{\varphi}(z) dz - 4 \sum_{k=2}^\infty \int_{-1/4}^{1/4} \hat{\varphi}(z+k) dz = \\ &= \varphi(0) - 4 \int_{z_1}^{z_2} \hat{\varphi}(z) dz - 4 \int_0^{1/4} \left(\sum_{k=-\infty}^\infty \hat{\varphi}(z+k) - \hat{\varphi}(z) - \hat{\varphi}(z-1) - \hat{\varphi}(z+1) \right) dz = \\ &= \varphi(0) - 4 \int_{z_1}^{z_2} \hat{\varphi}(z) dz - 4 \int_0^{1/4} \varphi(0) dz + 4 \left(\int_0^{1/4} + \int_{-1}^{-3/4} + \int_1^{5/4} \right) \hat{\varphi}(z) dz = \\ &= 4 \left(\int_0^{q_1} + \int_{q_2}^{q_3} - \int_{z_1}^{z_2} \right) \hat{\varphi}(z) dz. \end{aligned}$$

In these calculations we use parity $\sum_{k=-\infty}^\infty \hat{\varphi}(z+k) = \varphi(0)$; it following from Poisson summation formula $\sum_{\nu \in \mathbb{Z}} \hat{\varphi}(z+\nu) = \sum_{\nu \in \mathbb{Z}} \varphi(\nu) e(-\nu z)$ and $\varphi(\nu) = 0$ for $|\nu| \geq 1$. Theorem is proved.

REMARK 2. It is possible to improve new lower bound of the constant c , but it demands large calculations (these calculations are not included in this short article).

All computer calculations are obtained by mathematical package **Maple**.

REFERENCES

- [1] TAYKOV L.V. About the best approximation of Dirichlet kernels// Mathematical Notes 1993. V. 53. N 6. P. 116–121.
- [2] TAYKOV L.V. One range of extremum problems for trigonometric polynomials // UMN. 1965. V. 20. N 3. P. 205–211.
- [3] STEPANEC A.I. Uniform approximations by trigonometric polynomials. Kiev: Naukova Dumka, 1981.

E-mail address: `dvg@uic.tula.ru`

URL: `http://home.uic.tula.ru/~gd030473`